Earliest Age for Vaccine	Immunization	Notes
At birth	Hepatitis B Vaccine	Funded by OHIP in grade 7 but can be given earlier. Appears to be no decrease in protective effect if given at early age. Schedule is 0, 1, 6 months. Less immunogenicity reported if baby is < 2000 grams.
2 months	Group B Meningitis Vaccine Bexsero	As of Dec 2016, still not funded by OHIP. **For infants 2-5 months**, 4 doses should be given at 2, 4, and 6 months followed by a booster at 12-23 months. **For infants 6-11 months**, 3 doses should be given with at least 2 months separating them – third dose should be given in the second year of life. **For infants 12-23** months, only 2 doses are required at least 2 months apart. **For children 2-10**, 2 doses must be given at least 2 months apart. **For children 11-17,** 2 doses should be given at least 1 month apart.
	Group C Meningitis Vaccines	Funded by OHIP starting at 12 months and given again in schools during the grade 7 year. See next page for various Meningitis C vaccines available. For infants 2-12 months 3 doses at least 1 month

	Menjugate	apart followed by a booster dose (usually after 12 months). Infants 12 months and over need only a single dose (it is funded by OHIP at 12 months).
	NeisVac-C	**If less than 4 months,** baby receives 2 doses (0.5 ml) at least 2 months apart. Second dose must happen after baby is 5 months or older. **If baby is 4 months or older,** they can receive 1 dose only (0.5 ml).
	Group ACYW135 Meningitis Vaccines	Schedules exist for these vaccines starting at 2 months but they are not recommended by NACI for healthy children. If the child is high risk Menveo is the preferred ACYW135 vaccine at 2-23 months.
9 Years	HPV Vaccines (Gardasil, Gardasil 9, Cervarix)	Vaccine is not OHIP-covered until grade 8. Indicated for boys and girls ages 9-45 and boys ages 9-26. Safety and efficacy data not available outside of these ages.

*If you have any questions about the vaccines above, we would be happy to give you more information.